

Tsevegsuren, N., Davaakhuu, G., & Udval, T. (2014). Phytochemical analysis of *Cynara scolymus* L. cultivated in Mongolia. *Mongolian Journal of Chemistry*, **15**(41), 40–42.

Venugopal, K., Ahmad, H., Manikandan, E., Thanigai Arul, K., Kavitha, K., et al. (2017). The impact of anticancer activity upon *Beta vulgaris* extract mediated biosynthesized silver nanoparticles (ag-NPs) against human breast (MCF-7), lung (A549), and pharynx (Hep-2) cancer cell lines. *Journal of Photochemistry and Photobiology B: Biology*, **173**(January), 99–107.

Vongsak, B., Sithisarn, P., Mangmool, S., Thongpraditchote, S., Wongkrajang, Y., & Gritsanapan, W. (2013). The appropriate extraction method maximizes total phenolics, total flavonoid contents, and antioxidant activity of *Moringa oleifera* leaf extract. *Industrial Crops and Products*, **44**, 566–571.

Yang, M., Ma, Y., Wang, Z., Khan, A., Zhou, W., et al. (2020). Phenolic constituents, antioxidant and cytoprotective activities of crude extract, and fractions from cultivated artichoke inflorescence. *Industrial Crops and Products*, **143**(June 2019), 111433.

Zuorro, A., Maffei, G., & Lavecchia, R. (2016). Reuse potential of artichoke (*Cynara scolimus* L.) waste for the recovery of phenolic compounds and bioenergy. *Journal of Cleaner Production*, **111**(November 2017), 279–284.

Submitted: 25/06/2020

Revised: 09/04/2021

Accepted: 23/04/2021

DOI: 10.48129/kjs.10012