

Adaptive multi-step differential transformation method to solve ODE systems

AHMET GÖKDOĞAN* AND MEHMET MERDAN*

**Gümüşhane University, Department of Mathematical Engineering, 29100, Gümüşhane, Turkey*

ABSTRACT

In this paper, it is given a fast algorithm to solve chaotic differential systems using the multi-step differential transforms method (MsDTM). The approach is applied to a number of chaotic nonlinear differential equations and numerical results are given. Performance analyses reveal that the proposed approach is an efficiency tool to solve using fewer time step to the considered equation systems.

Keywords: The differential transform method; adaptive; Runge-Kutta method; chaotic systems.

INTRODUCTION

The differential transform method (DTM) is a semi-analytical-numerical method for solving integral equations, ordinary, partial differential equations and differential equation systems. The method enables the solution in terms of convergent series with easily computable components. The concept of the differential transform was first proposed by Zhou (1986) and its main application concern with both linear and nonlinear initial value problems in electrical circuit analysis. The DTM gives exact values of the n th derivative of an analytic function at a point in terms of known and unknown boundary conditions in a fast manner. This method generates, for differential equations, an analytical solution in the form of a polynomial. It is different from the traditional high order Taylor series method, which requires symbolic computations of the necessary derivatives of the data functions. The Taylor series method is computationally taken long time for large orders. The DTM is an iterative procedure for obtaining analytic Taylor series solutions of differential equations. Different applications of DTM can be found in Jang *et al.* (1997); Chen & Liu (1998); Ho & Chen (1998); Jordan & Smith (1999); Chen & Ho (1999a, 1999b); Jang *et al.* (2000a, 2000b, 2001); Köksal & Herdem (2002); Merdan *et al.* (2011a); Abdel-Halim Hassan (2002); Yildirim *et al.* (2012); Ayaz (2004); Arikoğlu & Özkol (2005); Merdan *et al.* (2011b); Ertürk & Momani (2007); Chang & Chang (2008); Ravi & Aruna (2008); Momani & Ertürk (2008);

El-Shahed (2008); Ravi & Aruna (2009a,2009b); Ebaid (2010); Merdan & Gökdoğan (2011).

However, DTM has some drawbacks. By using the DTM, we obtain a series solution, actually a truncated series solution. This series solution does not exhibit the real behaviors of the problem but gives a good approximation to the true solution in a very small region. To overcome the shortcoming, MsDTM was presented in Odibat *et al.* (2010); Gökdoğan *et al.* (2012a, 2012b). On the other hand, MsDTM has also some drawbacks. By using the DTM, the interval $[0, T]$ is divided into M sub-interval and the series solutions is obtained in $t \in [t_i, t_{i+1}]$, $i = 0, 1, \dots, M - 1$. In some problems, MsDTM can be required a very small sub-division of intervals. In this case, both the solution time lengthens and series solutions are obtained for a great number of sub-intervals.

In this study, we present the fast and effective algorithm to eliminate the above-mentioned disadvantages of classical MsDTM (Gökdoğan *et al.*, 2012a). The speed and efficiency of the provided algorithm were evaluated on the chaotic differential equations.

DIFFERENTIAL TRANSFORM METHOD

The basic definitions of differential transform are introduced as follows. Let $x(t)$ be analytic in a domain D and let $t = t_0$ represent any point in D . The function $x(t)$ is then represented by one power series whose center is located at t_0 . The differential transform of the k th derivative of a function $x(t)$ is defined as follows:

$$X(k) = \frac{1}{k!} \left[\frac{d^k x(t)}{dt^k} \right]_{t=t_0}, \forall t \in D \quad (1)$$

In (1), $x(t)$ is the original function and $X(k)$ is the transformed function. The differential inverse transformation of $X(k)$ is defined as follows:

$$x(t) = \sum_{k=0}^{\infty} X(k)(t - t_0)^k, \forall t \in D \quad (2)$$

From (1) and (2), we obtain

$$x(t) = \sum_{k=0}^{\infty} \frac{(t - t_0)^k}{k!} \left[\frac{d^k x(t)}{dt^k} \right]_{t=t_0}, \forall t \in D \quad (3)$$

The fundamental theorems of the one-dimensional differential transform are:

Theorem 1. If $z(t)=x(t)\pm y(t)$, then $Z(k)=X(k)\pm Y(k)$.

Theorem 2. If $z(t)=cy(t)$, then $Z(k)=cY(k)$.

Theorem 3. If $z(t) = \frac{dy(t)}{dt}$, then $Z(k)=(k + 1)Y(k + 1)$.

Theorem 4. If $z(t)=\frac{d^n y(t)}{dt^n}$, then $Z(k)=\frac{(k + n)!}{k!} Y(k)$.

Theorem 5. If $z(t) = x(t)y(t)$, then $Z(k) = \sum_{k_1=0}^k X(k_1)Y(k - k_1)$.

Theorem 6. If $z(t)=t^n$, then $Z(k) = \delta(k - n) = \begin{cases} 1 & k = n \\ 0 & k \neq n \end{cases}$.

In real applications, the function $x(t)$ is expressed by a finite series and (3) can be written as

$$x(t) = \sum_{k=0}^N X(k)(t - t_i)^k, \forall t \in D \tag{4}$$

Equation (4) implies that $\sum_{k=N+1}^{\infty} X(k)(t - t_i)^k$ is negligibly small. To illustrate the differential transform method (DTM) for solving differential equations systems, Consider a general system of first-order ODES

$$\begin{aligned} \frac{dx_1}{dt} + h_1(t, x_1, x_2, \dots, x_m) &= g_1(t), \\ \frac{dx_2}{dt} + h_2(t, x_1, x_2, \dots, x_m) &= g_2(t), \\ &\vdots \\ \frac{dx_m}{dt} + h_m(t, x_1, x_2, \dots, x_m) &= g_m(t), \end{aligned} \tag{5}$$

subject to the initial conditions

$$x_1(t_0) = d_1, \quad x_2(t_0) = d_2, \dots, x_m(t_0) = d_m. \tag{6}$$

According to DTM, by taking differential transformed both sides of the systems of equations given Eq. (5) and (6) is transformed as follows:

$$\begin{aligned} (k + 1)X_1(k + 1) + H_1(k) &= G_1(k), \\ (k + 1)X_2(k + 1) + H_2(k) &= G_2(k), \\ &\vdots \\ (k + 1)X_m(k + 1) + H_m(k) &= G_m(k). \end{aligned} \tag{7}$$

$$X_1(0) = d_1, \quad X_2(0) = d_2, \dots, X_m(0) = d_m. \quad (8)$$

Therefore, according to DTM the N -term approximations for (5) can be expressed as

$$\begin{aligned} \varphi_{1,n}(t) &= x_1(t) = \sum_{k=1}^N X_1(k)t^k, \\ \varphi_{2,n}(t) &= x_2(t) = \sum_{k=1}^N X_2(k)t^k, \\ &\vdots \\ \varphi_{m,n}(t) &= x_m(t) = \sum_{k=1}^N X_m(k)t^k. \end{aligned} \quad (9)$$

MULTI-STEP DIFFERENTIAL TRANSFORM METHOD

The approximate solutions (3) are generally, as will be shown in the numerical experiments of this paper, not valid for large t . A simple way of ensuring validity of the approximations for large t is to treat (9) as an algorithm for approximating the solutions of (4) in a sequence of intervals choosing the initial approximations as

$$\begin{aligned} x_{1,0}(t) &= x_1(t^*) = d_1^*, \\ x_{2,0}(t) &= x_2(t^*) = d_2^*, \\ &\vdots \\ x_{m,0}(t) &= x_m(t^*) = d_m^*. \end{aligned} \quad (10)$$

In order to carry out the iterations in every subinterval $[0, t_1)$, $[t_1, t_2)$, $[t_2, t_3)$, ..., $[t_{j-1}, t_j)$ of equal length h , we would need to know the values of the following (Odibat *et al.*, 2010),

$$x_{1,0}^*(t) = x_1(t^*), \quad x_{2,0}^*(t) = x_2(t^*), \quad \dots, \quad x_{m,0}^*(t) = x_m(t^*). \quad (11)$$

But, in general, we do not have these information at our clearance except at the initial point $t^* = t_0$. A simple way for obtaining the necessary values could be by means of the previous n -term approximations $\varphi_{1,n}, \varphi_{2,n}, \dots, \varphi_{m,n}$ of the preceding subinterval, i.e.,

$$x_{1,0}^* \cong \varphi_{1,n}(t^*), \quad x_{2,0}^* \cong \varphi_{2,n}(t^*), \quad \dots, \quad x_{m,0}^* \cong \varphi_{m,n}(t^*). \quad (12)$$

THE ALGORITHM

By sub-dividing interval $[0, T]$ into equilength M numbered sub-intervals, MsDTM method finds DTM solutions of these sub-ranges. But, it is necessary to select the step length of the small to solve some differential equations and this requires finding DTM solutions in more sub-intervals. Hence, the solution time prolongs in large values of T . A new approach is needed to overcome this problem. For this, we recommend a fast algorithm (Gökdoğan *et al.*, 2012a).

We apply the DTM to system (5) over the interval $[0, t_1]$, we will obtain the following approximate solution,

$$\begin{aligned}
 \varphi_{1,1}(t) = x_1(t) &= \sum_{k=1}^n X_1(k)t^k, \quad t \in [0, t_1] \\
 \varphi_{2,1}(t) = x_2(t) &= \sum_{k=1}^n X_2(k)t^k, \quad t \in [0, t_1] \\
 &\vdots \\
 \varphi_{m,1}(t) = x_m(t) &= \sum_{k=1}^n X_m(k)t^k, \quad t \in [0, t_1]
 \end{aligned} \tag{13}$$

using the initial conditions $x_k(0) = d_k, \quad k = 1, 2, \dots, m$. For $i \geq 2$, at each subinterval $[t_{i-1}, t_i]$, if $\max_{i=1, \dots, m} \left| \phi_{i,j-1}^N(t_{i-1}) - \phi_{i,j-1}^{N-1}(t_{i-1}) \right| \geq \varepsilon$, we will use the initial conditions $x_k(t_{i-1}) = x_{k-1}(t_{i-1}), \quad k = 1, 2, \dots, m$ and apply the DTM to Eq. (5) over the interval $[t_{i-1}, t_i]$, where t_0 in Eq. (1) is replaced by t_{i-1} . If $\max_{i=1, \dots, m} \left| \phi_{i,j-1}^N(t_{i-1}) - \phi_{i,j-1}^{N-1}(t_{i-1}) \right| < \varepsilon$, we get $x_k(t) = x_{k-1}(t), \quad k = 1, 2, \dots, m$, not apply the DTM to Eq. (5) over the this interval. The process is repeated and generates a sequence of approximate solutions $\phi_{k,i}(t), \quad i = 1, 2, \dots, K \leq M, \quad k = 1, 2, \dots, m$ for the solution $x_k(t), \quad k = 1, 2, \dots, m$.

NUMERICAL RESULTS

In this section, some chaotic systems are solved by using the proposed approach and MsDTM. The performance analyses are given for both methods. For performance analyses, only the operations up to the finding step of $x_m(t)$'s are taken, the elapsed time for graphic drawings are not included in the analysis. Analyses are carried out by MAPLE 7 software in a PC with a Pentium 1.6 GHz and 512 MB of RAM.

Example 1

The Genesio system, proposed by Genesio (Olek 1994) and Tesi (Genesio &

Tesi, 1992), is one of paradigms of chaos since it captures many features of chaotic systems. The dynamic equations of the system is given by

$$\begin{cases} x' = y \\ y' = z \\ z' = az + by + cx + dx^2 \end{cases} \quad (14)$$

when $a = -1.2, b = -2.92, c = -6, d = 1$, by taking with the initial conditions

$$x(0) = 0.2, y(0) = -0.3, z(0) = 0.1 \quad (15)$$

We will apply classic DTM and the adaptive MsDTM to nonlinear ordinary differential Eq. (14)-(15). Applying classic DTM for Eq.(14)- (15)

$$\begin{aligned} X(k+1) &= Y(k)k+1 \\ Y(k+1) &= Z(k)k+1 \\ Z(k+1) &= \frac{\left(aZ(k) + bY(k) + cX(k) + d\sum_{k_1=0}^k X(k_1)X(k-k_1) \right)}{k+1} \end{aligned} \quad (16)$$

$$X(0) = 0.2, Y(0) = -0.3, Z(0) = 0.1$$

By applying the multi-step DTM to (14)- (15), $X_i(n), Y_i(n)$ and $Z_i(n)$ satisfy the following recurrence relations for $n = 0, 1, 2, \dots, N-1$.

$$\begin{aligned} X_i(k+1) &= \frac{Y_i(k)}{k+1} \\ Y_i(k+1) &= \frac{Z_i(k)}{k+1} \\ Z_i(k+1) &= \frac{\left(aZ_i(k) + bY_i(k) + cX_i(k) + d\sum_{k_1=0}^k X_i(k_1)X_i(k-k_1) \right)}{k+1} \end{aligned} \quad (17)$$

$$X_0(0) = 0.2, X_i(0) = x_{i-1}(t_i), Y_0(0) = -0.3, Y_i(0) = y_{i-1}(t_i), Z_0(0)$$

$$= 0.1, Z_i(0) = z_{i-1}(t_i), i = 1, 2, \dots, K \leq M$$

The solutions obtained from the proposed approach are given below. The performance analyses are obtained by MsDTM and the approach summarized in Table 1. It is showed that the proposed approach is very fast and effective.

Table 1. Comparison processing time and time-step for $t \in [0, 20]$

5-term MsDTM($h = 0.1$)		5-term the proposed approach ($h = 0.1$)		
processing time	time-step		processing time	time-step
0.530s	200	$\varepsilon = 0.01$	0.290s	53
		$\varepsilon = 0.001$	0.370s	98
		$\varepsilon = 0.0001$	0.580s	149
MsDTM($h = 0.01$)		5-term the proposed approach ($h = 0.01$)		
Processing time	time-step		processing time	time-step
5.038s	2000	$\varepsilon = 0.01$	0.761s	47
		$\varepsilon = 0.001$	0.831s	75
		$\varepsilon = 0.0001$	0.951s	121
MsDTM($h = 0.001$)		5-term the proposed approach ($h = 0.001$)		
processing time	time-step		processing time	time-step
274.334s	20000	$\varepsilon = 0.01$	4.517s	46
		$\varepsilon = 0.001$	4.667s	73
		$\varepsilon = 0.0001$	6.119s	117




Fig. 1. Comparison 5-term MsDTM with $h = 0.1$ and RK4 with $h = 0.001$ (a1) $x(t)$, (b1) $y(t)$, (c1) $z(t)$ when $\varepsilon = 0.01$, (a2) $x(t)$, (b2) $y(t)$, (c2) $z(t)$ when $\varepsilon = 0.001$ and (a3) $x(t)$, (b3) $y(t)$, (c3) $z(t)$ when $\varepsilon = 0.0001$

Figure 1 show the approximate solutions for chaotic Genesis system obtained using the proposed approach with $h = 0.1$ and RK4 method with $h = 0.001$. We can observe that local changes obtained using the approach are in high adaptation with RK4 method for small value of ε . At the same time, we present the absolute errors between the 5-term MsDTM ($h=0.1$) solutions and the RK4($h=0.001$) solution in Table 2. It is showed that results obtained for small values of ε are compatible with RK4 even at large time step.

Table 2. Differences between 5-term MsDTM with $h = 0.1$ and RK4 with $h = 0.001$ for different ε values in $t \in [0, 20]$

t	$\Delta = MsDTM_{0.1} - RK4_{0.001} $ and $\varepsilon = 0.01$			$\Delta = MsDTM_{0.1} - RK4_{0.001} $ and $\varepsilon = 0.001$			$\Delta = MsDTM_{0.1} - RK4_{0.001} $ and $\varepsilon = 0.0001$		
	Δx	Δy	Δz	Δx	Δy	Δz	Δx	Δy	Δz
2	1.783e-03	3.201e-03	4.375e-03	2.528e-04	1.803e-04	8.008e-04	1.687e-05	2.155e-06	5.926e-05
4	6.136e-03	9.659e-04	2.151e-02	5.348e-04	3.965e-04	2.089e-03	2.436e-05	3.681e-05	1.033e-04
6	9.748e-03	1.345e-02	4.124e-02	4.297e-04	1.771e-03	2.304e-03	7.440e-06	1.224e-04	7.986e-05
8	4.148e-03	4.055e-02	3.675e-02	3.834e-04	3.497e-03	2.619e-04	5.435e-05	2.027e-04	1.098e-04
10	1.551e-02	6.128e-02	3.231e-02	2.350e-03	3.861e-03	6.403e-03	1.858e-04	1.635e-04	5.418e-04
12	5.412e-02	6.020e-02	1.768e-01	4.707e-03	1.291e-03	1.791e-02	2.835e-04	8.540e-05	1.183e-03
14	7.782e-02	2.191e-03	2.752e-01	6.366e-03	4.817e-03	2.330e-02	3.764e-04	4.683e-04	1.367e-03
16	9.427e-02	1.520e-01	2.834e-01	3.811e-03	1.823e-02	1.510e-02	6.664e-05	1.291e-03	5.866e-04
18	4.329e-02	2.406e-01	3.484e-01	3.318e-03	2.865e-02	3.047e-02	1.913e-04	2.043e-03	1.878e-03
20	9.535e-03	1.893e-01	2.367e-01	4.073e-03	2.370e-02	2.403e-02	4.558e-04	1.663e-03	1.363e-03




Fig. 2. Chaotic attractors for Genesis system with time span [0, 100] using adaptive MsDTM

In Figure 2, we reproduce the well-known chaotic attractors of the Genesis system using 5-term the proposed approach solutions with $h = 0.01$.

Example 2

Consider chaotic Coulet system (Coulet *et al.* 1979; Arnodo *et al.*, 1981)

$$\begin{cases} x' = y \\ y' = z \\ z' = az + by + cx + dx^3 \end{cases} \tag{18}$$

when $a = 0.8, b = -1.1, c = -0.45, d = -1$, by taking with the initial conditions

$$x(0) = 0.1, y(0) = 0.41, z(0) = 0.31 \tag{19}$$

We will apply classic DTM and the adaptive MsDTM to nonlinear ordinary differential Eq. (18)-(19). Applying classic DTM for Eq.(18)-(19)

$$\begin{aligned}
X(k+1) &= \frac{Y(k)}{k+1} \\
Y(k+1) &= \frac{Z(k)}{k+1} \\
Z(k+1) &= \frac{\left(aZ(k) + bY(k) + cX(k) + \right. \\
&\quad \left. d \sum_{k_2=0}^k \sum_{k_1=0}^{k_2} X(k_1)X(k_2 - k_1)X(k - k_2) \right)}{k+1}
\end{aligned} \tag{20}$$

$$X(0) = 0.1, \quad Y(0) = 0.41, \quad Z(0) = 0.31$$

By applying the multi-step DTM to Eq. (18)-(19), $X_i(n)$, $Y_i(n)$ and $Z_i(n)$ satisfy the following recurrence relations for $n = 0, 1, 2, \dots, N-1$.

$$\begin{aligned}
X_i(k+1) &= \frac{Y_i(k)}{k+1} \\
Y_i(k+1) &= \frac{Z_i(k)}{k+1} \\
Z_i(k+1) &= \frac{\left(aZ_i(k) + bY_i(k) + cX_i(k) \right. \\
&\quad \left. + d \sum_{k_2=0}^k \sum_{k_1=0}^{k_2} X_i(k_1)X_i(k_2 - k_1)X_i(k - k_2) \right)}{k+1}
\end{aligned} \tag{21}$$

$$X_0(0) = 0.1, \quad X_i(0) = x_{i-1}(t_i), \quad Y_0(0) = 0.41, \quad Y_i(0) = y_{i-1}(t_i), \quad Z_0(0)$$

$$= 0.31, \quad Z_i(0) = z_{i-1}(t_i), \quad i = 1, 2, \dots, K \leq M$$

The performance analyses are obtained by MsDTM and the approach summarized in Table 3. It is showed that the proposed approach is very fast and effective.

Table 3. Comparison processing time and time-step for $t \in [0, 20]$

5-term MsDTM(h=0.1)		5-term the proposed approach MsDTM(h=0.1)		
processing time	time-step		processing time	time-step
1.262s	200	$\varepsilon = 0.01$	0.491s	48
		$\varepsilon = 0.001$	1.002s	137
		$\varepsilon = 0.0001$	1.352s	199
MsDTM(h=0.01)		5-term the proposed approach (h=0.01)		
processing time	time-step		processing time	time-step
14.421s	2000	$\varepsilon = 0.01$	0.911s	43
		$\varepsilon = 0.001$	1.021s	102
		$\varepsilon = 0.0001$	1.943s	228
MsDTM(h=0.001)		5-term the proposed approach (h=0.001)		
processing time	time-step		processing time	time-step
614.233s	20000	$\varepsilon = 0.01$	2.974s	43
		$\varepsilon = 0.001$	3.244s	100
		$\varepsilon = 0.0001$	4.066s	216




Fig. 3. Comparison MsDTM with $h = 0.1$ and $RK4$ with $h = 0.001$ (a1) $x(t)$, (b1) $y(t)$, (c1) $z(t)$ when $\varepsilon = 0.01$, (a2) $x(t)$, (b2) $y(t)$, (c2) $z(t)$ when $\varepsilon = 0.001$ and (a3) $x(t)$, (b3) $y(t)$, (c3) $z(t)$ when $\varepsilon = 0.0001$

Figure 3 show the approximate solutions for chaotic Coulet system obtained using adaptive MsDTM with $h = 0.1$ and RK4 method with $h = 0.001$. We can observe that local changes obtained using the approach are in adaptation with RK4 method for small value of ε .

Table 4. Differences between 5-term MsDTM with $h = 0.1$ and RK4 with $h = 0.001$ for different ε values in $t \in [0, 20]$

t	$\Delta = MsDTM_{0.1} - RK4_{0.001} $ and $\varepsilon = 0.01$			$\Delta = MsDTM_{0.1} - RK4_{0.001} $ and $\varepsilon = 0.001$			$\Delta = MsDTM_{0.1} - RK4_{0.001} $ and $\varepsilon = 0.0001$		
	Δx	Δy	Δz	Δx	Δy	Δz	Δx	Δy	Δz
2	2.776e-03	4.529e-03	3.159e-03	3.266e-04	2.717e-04	6.930e-04	4.255e-05	2.537e-05	9.442e-05
4	1.099e-02	9.492e-03	8.943e-03	8.381e-04	7.566e-04	6.223e-04	1.469e-04	9.007e-05	2.441e-04
6	1.658e-02	8.172e-03	5.696e-03	1.396e-03	3.834e-04	3.726e-04	1.844e-04	3.928e-05	6.347e-05
8	2.634e-02	1.693e-02	4.631e-02	1.677e-03	1.161e-03	2.747e-03	2.127e-04	1.406e-04	4.287e-04
10	6.142e-02	2.859e-02	5.362e-02	4.058e-03	1.879e-03	3.522e-03	5.604e-04	1.948e-04	6.672e-04
12	7.266e-02	7.440e-03	2.320e-02	4.782e-03	4.977e-04	1.269e-03	6.558e-05	2.993e-05	2.845e-04
14	1.504e-01	8.325e-02	5.004e-02	9.341e-03	5.031e-03	3.078e-03	2.163e-04	1.484e-04	3.850e-05
16	4.331e-01	2.053e-01	5.878e-02	2.581e-02	1.034e-02	5.794e-04	5.004e-04	1.863e-04	3.047e-05
18	8.356e-01	7.264e-02	2.990e-01	2.265e-02	2.801e-02	3.931e-02	4.521e-04	5.983e-04	8.671e-04
20	2.422e-01	5.472e-01	3.564e-02	5.059e-02	1.385e-03	7.466e-02	1.102e-03	2.455e-04	1.599e-03

we submit the absolute errors between the 5-term MsDTM ($h=0.1$) solutions and the RK4($h=0.001$) solution in Table 4. It is showed that results obtained for small values of ε are compatible with RK4 even at large time step.




Fig. 4. Chaotic attractors for Coulett system with time span $[0, 100]$ using adaptive MsDTM

In Figure 4, we reproduce the well-known chaotic attractors of the Coulett system using 5-term the approach solutions with $h = 0.01$.

Example 3

Here, we take Lorenz system into consideration. In 1963, the Lorenz system is introduced by Edward Lorenz, (1963). The nonlinear differential equations that describe the Lorenz system are

$$\begin{cases} x' = \rho(y - x) \\ y' = Rx - y - xz \\ z' = xy + bz \end{cases} \quad (22)$$

when $\rho = 10, b = -8/3, R = 28$, by taking with the initial conditions

$$x(0) = -15.8, y(0) = -17.48, z(0) = 35.64 \quad (23)$$

We will apply classic DTM and the adaptive MsDTM to nonlinear ordinary

differential Eq. (22)-(23). Applying classic DTM for Eq. (22)-(23)

$$\begin{aligned}
 X(k+1) &= \frac{\rho(Y(k) - X(k))}{k+1} \\
 Y(k+1) &= \frac{\left(RX(k) - Y(k) - \sum_{k_1=0}^k X(k_1)Z(k-k_1) \right)}{k+1} \\
 Z(k+1) &= \frac{\left(bZ(k) + \sum_{k_1=0}^k X(k_1)Y(k-k_1) \right)}{k+1}
 \end{aligned} \tag{24}$$

$$X(0) = 15.8, \quad Y(0) = -17.48, \quad Z(0) = 35.64$$

By applying the multi-step DTM to Eq. (22)-(23), $X_i(n)$, $Y_i(n)$ and $Z_i(n)$ satisfy the following recurrence relations for $n = 0, 1, 2, \dots, N-1$.

$$\begin{aligned}
 X_i(k+1) &= \frac{\rho(Y_i(k) - X_i(k))}{k+1} \\
 Y_i(k+1) &= \frac{\left(RX_i(k) - Y_i(k) - \sum_{k_1=0}^k X_i(k_1)Z_i(k-k_1) \right)}{k+1} \\
 Z_i(k+1) &= \frac{\left(bZ_i(k) + \sum_{k_1=0}^k X_i(k_1)Y_i(k-k_1) \right)}{k+1}
 \end{aligned} \tag{25}$$

$$X_0(0) = 15.8, \quad X_i(0) = x_{i-1}(t_i), \quad Y_0(0) = -17.48, \quad Y_i(0) = y_{i-1}(t_i), \quad Z_0(0)$$

$$= 35.64, \quad Z_i(0) = z_{i-1}(t_i), \quad i = 1, 2, \dots, K \leq M$$

The performance analyses are obtained by MsDTM and the approach summarized in Table 5. It is showed that the proposed approach is very fast and effective.

Table 5. Comparison processing time and time-step for $t \in [0, 10]$

5-term MsDTM($h = 0.1$)		5-term the proposed approach MsDTM($h = 0.1$)	
processing time	time-step	processing time	time-step
0.641s	100	$\varepsilon = 0.01$	0.471s
		$\varepsilon = 0.001$	0.480s
		$\varepsilon = 0.0001$	0.480s
MsDTM($h = 0.01$)		Adaptive MsDTM($h = 0.01$)	
processing time	time-step	processing time	time-step
6.529s	1000	$\varepsilon = 0.01$	0.981s
		$\varepsilon = 0.001$	1.112s
		$\varepsilon = 0.0001$	1.282s
MsTM($h = 0.001$)		Adaptive MsDTM($h = 0.001$)	
processing time	time-step	processing time	time-step
315.785s	10000	$\varepsilon = 0.01$	5.778s
		$\varepsilon = 0.001$	5.888s
		$\varepsilon = 0.0001$	6.159s




Fig. 5. Comparison 10-term MsDTM with $h = 0.1$ and RK4 with $h = 0.001$ (a1) $x(t)$, (b1) $y(t)$, (c1) $z(t)$ when $\varepsilon = 0.01$, (a2) $x(t)$, (b2) $y(t)$, (c2) $z(t)$ when $\varepsilon = 0.001$ and (a3) $x(t)$, (b3) $y(t)$, (c3) $z(t)$ when $\varepsilon = 0.0001$

Figure 5 show the approximate solutions for chaotic Lorenz system obtained using the approach with $h = 0.1$ and RK4 method with $h = 0.001$. We can observe that local changes obtained using the approach are in high adaptation with RK4 method for small value of ε .

Table 6. Differences between 10-term MsDTM with $h = 0.01$ and RK4 with $h = 0.001$ for different ε values in $t \in [0, 10]$

t	$\Delta = MsDTM_{0.1} - RK4_{0.001} $ and $\varepsilon = 0.01$			$\Delta = MsDTM_{0.1} - RK4_{0.001} $ and $\varepsilon = 0.001$			$\Delta = MsDTM_{0.1} - RK4_{0.001} $ and $\varepsilon = 0.0001$		
	Δx	Δy	Δz	Δx	Δy	Δz	Δx	Δy	Δz
2	7.490e-04	1.758e-03	4.903e-03	7.820e-05	3.861e-04	1.072e-04	3.350e-07	9.623e-06	1.529e-05
4	4.153e-04	2.449e-03	1.098e-02	4.234e-04	7.867e-04	2.777e-03	2.031e-05	1.308e-05	5.638e-05
6	2.128e-03	4.930e-03	9.280e-03	9.715e-04	3.165e-03	8.144e-04	2.646e-05	3.327e-04	2.834e-04
8	1.515e-03	5.008e-03	8.055e-03	7.427e-03	3.554e-03	1.501e-02	5.309e-04	2.024e-04	1.023e-03
10	2.150e-02	3.501e-02	9.312e-03	1.113e-02	1.817e-02	2.074e-03	6.323e-04	1.024e-03	3.757e-05
12	1.472e-01	2.477e-01	2.164e-02	8.109e-02	1.327e-01	2.128e-02	4.511e-03	7.378e-03	1.257e-03
14	1.446e-01	1.590e-01	1.055e-01	7.958e-02	7.959e-02	6.547e-02	4.358e-03	4.408e-03	3.521e-03
16	3.952e-01	6.251e-01	2.349e-01	1.882e-01	2.996e-01	1.117e-01	1.071e-02	1.713e-02	6.000e-03
18	6.305e+00	1.204e+01	2.465e+00	2.971e+00	5.794e+00	2.391e-01	1.611e-01	3.140e-01	2.897e-02
20	1.770e+01	1.188e+01	2.340e+00	2.106e+01	1.542e+01	7.081e-01	4.305e-02	8.495e-01	7.908e-01

we present the absolute errors between the 10-term MsDTM ($h=0.1$) solutions and the RK4($h=0.001$) solution in Table 6. It is showed that results obtained for small values of ε are compatible with RK4.




Fig. 6. Chaotic attractors for Lorenz system with time span $[0, 100]$ using adaptive MsDTM

In Figure 6, we reproduce the well-known chaotic attractors of the Genesio system using 5-term the approach solutions with $h = 0.01$.

CONCLUSION

In this work, we suggested a new efficient and fast algorithm for the MsDTM, a reliable modification of the DTM that develops the convergence of the series solution. Comparisons between the proposed method solutions and RK4's numerical solutions were made. The method provides immediate and visible symbolic terms of analytic solutions. The validity of the proposed method has been successfully demonstrated by applying it for the well-known chaotic systems (Genesio, Coulet, Lorenz). The results show that adaptive MsDTM provides the user remarkable performance in terms of processing time and time step.

ACKNOWLEDGEMENT

Authors thank to TÜBİTAK (The Scientific and Technological Research Council of TURKEY) for their financial support.

REFERENCES

- Abdel-Halim Hassan, I. H. 2002.** On solving some eigenvalue problems by using a differential Transformation. *Applied Mathematics and Computation***127**: 1-22.
- Arikoğlu, A. & Özkol, I. 2005.** Solution of boundary value problems for integro-differential equations by using differential transform method. *Applied Mathematics and Computation* **168**: 1145-1158.
- Arnodo, A., Couillet, P. & Tresser, C. 1981.** Possible new strange attractors with spiral structure. *Communications in Mathematical Physics* **79(4)**:573-579.
- Ayaz, F. 2004.** Solutions of the system of differential equations by differential transform Method. *Applied Mathematics and Computation* **147**: 547-567.
- Chang, S. H. & Chang, I. L. 2008.** A new algorithm for calculating one-dimensional differential transform of nonlinear functions. *Applied Mathematics and Computation***195**: 799-808.
- Chen, C. L. & Liu, Y.C. 1998.** Solution of two point boundary value problems using the differential transformation method. *Journal of Optimization Theory and Applications* **99**: 23-35.
- Chen, C. K. & Ho, S. H. 1999a.** Transverse vibration of a rotating twisted Timoshenko beam under axial loading using differential transform. *International journal of mechanical sciences* **41**: 1339-1356.
- Chen, C. K. & Ho, S. H. 1999b.** Solving partial differential equations by two-dimensional differential transform method. *Applied Mathematics and Computation* **106**: 171-179.
- Couillet, P., Tresser, C. & Arnodo, A. 1979.** Transition to stochasticity for a class of forced oscillators. *Physics Letters A* **72**:268-270.
- Ebaid, A. E. H. 2010.** Approximate periodic solutions for the non-linear relativistic harmonic oscillator via differential transformation method. *Communications in Nonlinear Science and Numerical Simulation* **15**: 1921-1927.
- El-Shahed, M. 2008.** Application of differential transform method to non-linear oscillatory Systems. *Communications in Nonlinear Science and Numerical Simulation* **13**: 1714-1720.
- Ertürk, V. S. & Momani, S. 2007.** Comparing numerical methods for solving fourth-order boundary value problems. *Applied Mathematics and Computation* **188 (2)**: 1963- 1968.
- Genesio, R. & Tesi, A. 1992.** Harmonic balance methods for the analysis of chaotic dynamics in nonlinear systems. *Automatica* **28**: 531-548.

- Gökdoğan, A., Merdan, M. & Yildirim, A. 2012a.** Adaptive multi-step differential transformation method for solving nonlinear differential equations. *Mathematical and Computer Modelling* **55**: 761-769.
- Gökdoğan, A., Merdan, M. & Yildirim, A. 2012b.** A multistage differential transformation method for approximate solution of Hantavirusinfection model. *Communications in Nonlinear Science and Numerical Simulation* **17(1)**: 1-8.
- Ho, S. H. & Chen, C. K. 1998.** Analysis of general elastically end restrained non-uniform beams using differential transform. *Applied Mathematical Modelling* **22**: 219-234.
- Jang, M. J., Chen, C. L. & Liu, Y. C. 1997.** Analysis of the response of a strongly nonlinear damped system using a differential transformation technique. *Applied Mathematics and Computation* **88**: 137-151.
- Jang, M. J., Chen, C. L. & Liy, Y. C. 2000a.** On solving the initial-value problems using the differential transformation method. *Applied Mathematics and Computation* **115**: 145-160.
- Jang, M. J., Chen, C. L. & Liu, Y. C. 2000b.** On solving the initial value problems using the differential transformation method. *Applied Mathematics and Computation* **115**: 145-160.
- Jang, M. J., Chen, C. L. & Liu, Y. C. 2001.** Two-dimensional differential transform for partial differential equations. *Applied Mathematics and Computation* **121**: 261-70.
- Jordan, D. W. & Smith, P. 1999.** *Nonlinear Ordinary Differential Equations.* Oxford University Press.
- Köksal, M. & Herdem, S. 2002.** Analysis of nonlinear circuits by using differential Taylor Transform. *Computers & Electrical Engineering* **28**: 513-525.
- Lorenz, E. 1963.** Deterministic nonperiodic flow. *Journal of the Atmospheric Sciences* **20**: 130-141.
- Merdan, M. & Gökdoğan, A. 2011.** Solution Of Nonlinear Oscillators With Fractional Nonlinearities By Using The Modified Differential Transformation Method. *Mathematical and Computational Applications* **16(3)**: 761-772.
- Merdan, M., Gökdoğan, A & Ertürk, V. S. 2011a.** On Solving Coulet System by Differential Transformation Method. *Çankaya University Journal of Science and Engineering* **8(1)**: 111-121.
- Merdan, M., Gökdoğan, A & Ertürk, V. S. 2011b.** A numeric-analytic method for approximating three-species food chain models. *International Journal of the Physical Sciences* **6(2)**: 1822-1833.

- Momani, S. & Ertürk, V. S. 2008.** Solutions of non-linear oscillators by the modified differential transform method. *Computers & Mathematics with Applications* **55(4)**: 833- 842.
- Odibat, Z. M., Bertelle, C., Aziz-Alaoui, M. A. & Duchamp, G. H. E. 2010.** A multi-step differential transform method and application to non-chaotic or chaotic systems. *Computers & Mathematics with Applications* **59**:1462-1472.
- Olek, S. 1994.** An accurate solution to the multispecies Genesio equations. *SIAM Review* **36(3)**: 480-488.
- Ravi, A. S. V. K. & Aruna, K. 2008.** Solution of singular two-point boundary value problems using differential transformation method. *Physics Letters A* **372**: 4671-4673.
- Ravi, A. S. V. K. & Aruna, K. 2009a.** Two-dimensional differential transform method for solving linear and non-linear Schrödinger equations. *Chaos Solitons & Fractals* **41**: 2277-2281.
- Ravi, A. S. V. K. & Aruna, K. 2009b.** Differential transform method for solving the linear and nonlinear Klein-Gordon equation. *Computer Physics Communications* **180**: 708-711.
- Yildirim, A., Merdan, M. & Gökdoğan, A. 2012.** Chaotic systems via multi-step differential transformation method. *Canadian Journal of Physics* **90**:1-16.
- Zhou, J. K. 1986.** *Differential Transformation and its Applications for Electrical Circuits*, Huazhong University Press, Wuhan, China.

Submitted : 3/10/2011

Revised : 20/12/2011

Accepted : 1/5/2012

طريقة تحويل تفاضلي تكيفي متعدد الخطوات لحل أنظمة المعادلات التفاضلية الاعتيادية

**أحمد جوكدوجان ، **محمد مردان

قسم الهندسة الرياضية - جامعة جوموشان 29100 - تركيا

خلاصة

نقدم في هذا البحث خوارزمية سريعة لحل الأنظمة التفاضلية الشواشية وذلك باستخدام طريقة تحويل تفاضلي تكيفي متعدد الخطوات. ونطبق هذه الخوارزمية على عدد من المعادلات التفاضلية الشواشية وغير الخطية فنحصل على نتائج عددية. ويكشف تحليل الأداء أن طريقتنا المقترحة ذات فعالية وتحتاج إلى وقت وخطوات أقل لحل الأنظمة موضوع الدراسة.

مجلة فصلية أكاديمية

محكمة تعنى بنشر البحوث

والدراسات القانونية والشرعية

تصدر عن مجلس النشر العلمي - جامعة الكويت

مجلة الحقوق



رئيس التحرير

الأستاذ الدكتور أحمد عبد الرحمن الملحم



صدر العدد الأول في

يناير ١٩٧٧

الاشتراكات

في الكويت	في الدول العربية	في الدول الأجنبية
٣ دولارات	٤ دولارات	١٥ دولاراً
١٥ ديناراً	١٥ ديناراً	٦٠ دولاراً

المراسلات

توجه جميع المراسلات إلى رئيس التحرير على العنوان الآتي:

مجلة الحقوق - جامعة الكويت ص.ب: ٦٤٩٨٥ الشويخ - ب 70460 الكويت

تلفون: ٢٤٨٣٥٧٨٩ - ٢٤٨٤٧٨١٤ فاكس: ٢٤٨٣١١٤٢

E.mail: jol@ku.edu.kw

عنوان المجلة في شبكة الإنترنت <http://www.pubcouncil.kuniv.edu.kw/jol>

ISSN 1029 - 6069